

## Streamlining Various Definitions of Childhood Obesity

Syed Arif Kamal<sup>†</sup>

SF Growth-and-Imaging Laboratory, the NGDS Pilot Project, University of Karachi, Karachi 75270, Pakistan; e-mail: [profdrakamal@gmail.com](mailto:profdrakamal@gmail.com)

Obesity has become a universal problem among children. Childhood obesity may be connected to grave psychological, physical and social consequences. First lady of United States, **Her Excellency, Michelle Obama** declared childhood obesity an epidemic for her country. Obesity develops when there exists a discrepancy between intake and output of energy, disturbing the original steady state and formation of a fresh steady state at a higher level, resulting in increased body-fat storage. In a 2011 work, the author described steady state as a situation employing transfer of energy at a uniform rate (energy-transfer perspective). This is a condition in which probability of occupation is not same in different states; however, it does not vary with time (probability-of-occupation perspective). The delicate balance between tissue synthesis (gain of height) and fat storage (obesity), if mathematically modeled, may prevent obesity. According to Poskitt (1995), representing ECOG, lack of childhood-obesity definition has been a matter of concern for the group. In 2000 she mentioned that the concept of relative *BMI* had been generally accepted despite considerable imprecision in defining obesity. In 2001 she stated that *BMI* could not be considered as offering the ‘best’ definition, although it might be ‘useful’ and ‘practical’ for epidemiological, clinical and population-research purposes. She, further, added that work on definition was essential and needed continuing reassessment. Cole *et al.* (2000) defined childhood obesity based on pooled-international data and linked to adult-obesity-cutoff point of *BMI* to be  $30 \text{ kg/m}^2$ . Flegal *et al.* (2010) divided *BMI*-for-age categories into three ranges: ‘normal’, ‘intermediate’ and ‘high’. The first one *most unlikely*, whereas the last one *most likely*, to have high adiposity. Rolland-Cachera *et al.* (2011), on behalf of ECOG, defined three main cutoffs of *BMI*, constituting four ranges: ‘thin’, ‘normal’, ‘overweight’ and ‘obese’. Zhao and Grant (2011) observed that obesity might be defined as excess of body fat. Al-Gindan *et al.* (2015) are of the opinion that most national-survey analyses equating *BMI* in excess of  $30 \text{ kg/m}^2$  with ‘obesity’ lead to survey-data misinterpretation. The author is of the opinion that ‘overweight’ (classified based on measurement of ‘net mass’ — mass obtained without any clothing) needs to be distinguished from ‘over-fat’ (classified based on measurement of skinfolds as well as waist and hip circumferences). One needs a definition based, solely, on measurement of mass, not measurement of fat, which is difficult to obtain in a reproducible manner. Starting from this year, mass can be recorded to least count of  $0.005 \text{ kg}$  in SF Growth-and-Imaging Laboratory. Kamal and Razzaq (2014) investigated reproducibility of mass measurement to least count of  $0.01 \text{ kg}$ . The author defines childhood obesity as the condition in which a youngster is required to shed off net mass at the end of 6-month period as compared to current mass based on ‘Growth-and-Obesity Vector-Roadmap’ recommendations. In a forthcoming paper, ‘In Search of a Definition of Childhood Obesity’, the author intends to compare ‘*BMI*-based-optimal mass’ with ‘height-percentile-based-optimal mass’ and give mathematical relationship for losing net mass within 6 months.

### Already known on this topic

- Childhood obesity a serious public health concern

---

- Obesity a complex disease that involves interactions between environmental and genetic factors

---

- The true prevalence of childhood obesity difficult to empirically quantify as there is currently no internationally-accepted definition

---

- BMI* still the most popular method of classifying fatness and thinness

---

- Various definitions obesity proposed included relative *BMI*, cutoff point as  $30 \text{ kg/m}^2$  (adult *BMI*), *BMI* ranges (below 85<sup>th</sup> percentile: normal, 85<sup>th</sup> to 95<sup>th</sup> percentile: intermediate, equal to or above 95<sup>th</sup> percentile: high)

### SF Growth-and-Imaging Laboratory contributions

- 2004 Optimal mass (mention of name; formal definition in 2011)

---

- 2011 Statuses (pertaining-to-mass) and (pertaining-to-height)

---

- 2012 Estimated-adult *BMI*

---

- 2013-2016 1<sup>st</sup>- to 4<sup>th</sup>-generation solutions of childhood obesity

---

- 2014 Energy-channelization I-III, which included puberty-induced energy-channelization

---

- 2014 Pseudo-gain of mass and height

---

- 2014 Use of height- and mass-percentile trajectories instead of growth (height) velocity and rate of mass gain/loss

---

- 2014 CDC Growth Tables extended to include percentiles in the range 0.01<sup>th</sup> to 99.99<sup>th</sup> (to handle extreme cases)

---

- 2015 Month-wise targets (next 6 months) to shed-off mass

---

- 2016 Mass and height measurements to least counts of  $0.005 \text{ kg}$  and  $0.005 \text{ cm}$ , respectively, accompanied by manual, version 9.1

### This work adds

Definition of childhood obesity — A child is considered obese if the incumbent is required to lose net mass (weight) within the next 6 months

### The next step

- Mathematical relationship among percentiles of mass, height, mid-parental height and army-cutoff height that warrants losing net mass (weight) within the next 6 months

---

- Investigation of suitability of *BMI* use as childhood-obesity indicator

---

- 5<sup>th</sup>-generation solution of childhood obesity

September 4, 2013	1 <sup>st</sup> -Generation Sol. <a href="http://www.ngds-ku.org/Presentations/Roadmap.pdf">http://www.ngds-ku.org/Presentations/Roadmap.pdf</a>
September 4, 2014	2 <sup>nd</sup> -Generation Sol. <a href="http://www.ngds-ku.org/Presentations/Enhanced.pdf">http://www.ngds-ku.org/Presentations/Enhanced.pdf</a>
June 1, 2015	3 <sup>rd</sup> -Generation Sol. <a href="http://www.ngds-ku.org/Papers/J38.pdf">http://www.ngds-ku.org/Papers/J38.pdf</a>
February 13, 2016	4 <sup>th</sup> -Generation Sol. <a href="http://www.ngds-ku.org/Presentations/Vector.pdf">http://www.ngds-ku.org/Presentations/Vector.pdf</a>
January 1, 2017	5 <sup>th</sup> -Generation Sol. <a href="http://www.ngds-ku.org/Papers/J45.pdf">http://www.ngds-ku.org/Papers/J45.pdf</a> (expected)



### Solutions of childhood-obesity problem proposed in SF Growth-and-Imaging Laboratory, University of Karachi

**Keywords:** Body-mass index (*BMI*), estimated-adult *BMI*, optimal mass, Growth-and-Obesity Vector-Roadmap, ECOG (European Childhood Obesity Group)

**Web address of this document:** <http://www.ngds-ku.org/Presentations/Childhood-Obesity-Definitions.pdf>

**HTML version:** <http://www.ngds-ku.org/pub/confabstB.htm#C131>:

<sup>†</sup>Prof. Dr. Syed Arif Kamal; PhD (Mathematical Neuroscience); MA, Johns Hopkins, Baltimore, MD, United States; Ex-Member, American Alliance for Health, Physical Education, Recreation and Dance (AAHPERD), Ex-Chairman, Department of Health, Physical Education and Sports Sciences; Ex-Dean, Faculties of Science and Engineering; Ex-Acting Vice Chancellor, University of Karachi • *paper mail*: Project Director, the NGDS Pilot Project, PO Box 8423, Karachi 75270, Pakistan • *homepage*: <http://www.ngds-ku.org/kamal> • *project URL*: <http://ngds-ku.org>