

## Effective Decision Making for Presence of Scoliosis Based on Moiré Fringe Topography

Syed Arif Kamal<sup>¶</sup>, Maqsood Sarwar<sup>§</sup> and Ani Haider<sup>§</sup>

<sup>¶</sup> <sup>§</sup>SF Growth-and-Imaging Laboratory, Anthromathematics Group, Department of Mathematics, University of Karachi, Karachi 75270, Pakistan; <sup>¶</sup>Subject Committee for Physical Education, Health and Sport Sciences, National Testing Service Pakistan; <sup>¶</sup>[profdrakamal@gmail.com](mailto:profdrakamal@gmail.com)

Visual (⊥) & Forward Bending (⊥) POSITIVE Hip-Weakness Suspected (Tredelenburg Sign)	Visual (↑) & Forward Bending (↑) POSITIVE Leg-Length-Inequality Suspected (Spinal Dimples)
Visual (⊥) & Visual (↑) POSITIVE Postural-Problem Suspected (Mild-Stretching Exam)	Forward Bending (⊥) & Forward Bending (↑) POSITIVE Spinal-Rotation Suspected (Moiré Fringe Topography)

**Fig. 1. Decision matrix indicating presence of spinal rotation: test conducted in sitting (⊥)/standing (↑) position**

Scoliosis, lateral curvatures and rotations of the spinal column, is a body-disfiguring condition. A two-minute-stripped-orthopedic examination may be able to alert the physician to early-warning signs. Our group tested a protocol in a local school, which was applied to seven- and eight-year-old students. Figure shows decision matrix to detect possible presence of spinal rotation. This matrix is based on four tests, visual (standing), visual (sitting), forward bending (standing) and forward bending (sitting) — postural problem suspected through positive visual examinations (standing and sitting), indicated through positive mid-stretching test; leg-length in-equality suspected though positive visual and forward-bending tests (both standing), indicated through uneven spinal dimples; hip weakness suspected through positive visual and forward-bending tests (both sitting), indicated through positive Tredelenburg sign; spinal rotation suspected through positive forward-bending tests (standing and sitting), indicated through positive moiré.


positive visual and forward-bending tests (both sitting), indicated through positive Tredelenburg sign; spinal rotation suspected through positive forward-bending tests (standing and sitting), indicated through positive moiré.

**Keywords:** Scoliosis risk, visual examination, forward-bending test, tall children, wasted children

**Conflict of Interest Statement:** No potential conflict of interest is identified for this work

**Grant Sponsor:** Dean's (Science) Research Grant, University of Karachi, number DFSR/2009

**Research Ethics:** Project initiated after Institutional Review Process and conducted in compliance with ethical and human-right standards in our region.



**Fig. 2a-f. Early-warning signals for presence of scoliosis, visual — (a) front, (b) side and (c) back and (d) forward bending checks as well as moiré topographs of (e) front and (f) back**

Web address of this document: <http://www.ngds-ku.org/Presentations/Decision.pdf>

HTML version: <http://www.ngds-ku.org/pub/confabstB.htm#C117>:

<sup>¶</sup>PhD (Mathematical Neuroscience); MA, Johns Hopkins, Baltimore, MD, United States; MS, Indiana, Bloomington, IN, United States; Project Director, the NGDS Pilot Project; Director, SF Growth-and-Imaging Laboratory; Head, Anthromathematics Group, University of Karachi; Member, Subject Committee for Physical Education, Health and Sport Sciences, National Testing Service Pakistan; Sessional faculty, the Aga Khan University Medical College (1996-2006); Associated Professor in Orthopedic Surgery, Malmö General Hospital, Sweden (1988); Research Associate in Orthopedic Surgery, James Whitecomb Riley Hospital for Children, Indianapolis, IN, United States (1980) and; *paper mail*: Professor and Chairman, Department of Mathematics, University of Karachi, Karachi 75270, Pakistan; *telephone*: +92 21 9926 1300-15 ext. 2293; *homepage*: <http://www.ngds-ku.org/kamal>; *project URL*: <http://ngds-ku.org>