

Paper presented during *the Second Conference on Athromathematics and Sport Mathematics in the Memory of (Late) Hussain Ahmed Bilgirami (ANTHROMATHEMATICS 2014)*, Department of Mathematics, University of Karachi, Karachi, Pakistan and Government College, Hyderabad, Pakistan, Thursday, September 4, 2014 (1315h-1330h, Assembly Hall, Govt. College), Technical Session Anthro14-II, abstract#Anthro14-06, p 10

## Crystal-Structure-Concept-Based Modeling of the Human Spinal Column Validated through 3-D-Bone Scanning

Syed Arif Kamal<sup>†\*</sup>, Maqsood Sarwar<sup>†</sup> and Muhammed Khalid Rajput<sup>§</sup>

SF Growth-and-Imaging Laboratory, Anthromathematics Group, Departments of <sup>†</sup>Mathematics and <sup>§</sup>Physics, University of Karachi, Karachi 75270, Pakistan; <sup>†</sup>[profdrakamal@gmail.com](mailto:profdrakamal@gmail.com)

Spinal column, being a 3-D structure, is not completely visualized in ordinary anteroposterior (AP) X rays, which show spinal projection in the frontal plane. Hence, such X rays do not display kyphosis or lordosis. 3-D-spinal models were able to generate full views from projections of spine in the frontal and the sagittal planes obtained from AP- and lateral-X-ray pictures or moiré topographs of patients in the attention position. A parabolic curve was generated from moiré measurements,  $x = x(\xi)$ ,  $y = y(\xi)$ ,  $z = z(\xi)$ , best fitted to discrete measurements performed at locations of 33 vertebrae of spinal column,  $\xi_i$ ;  $i = 1, \dots, 33$  (cervical, thoracic, lumbar and sacral regions). Rotation of coordinate system through a suitable angle made the cross term ( $yz$ ) vanish and represented curvatures as double the coefficients of squares of  $y$  and  $z$ . These curvatures were used to define degree of correction of spinal deformity (<http://www.ngds-ku.org/Papers/J18.pdf>). This 3-D-static model was used to study children's postures. The dynamic model was constructed by including spinal-column movements during a gait cycle, linking spinal column in the anatomical position to the first phase of gait cycle <http://www.ngds-ku.org/Papers/J16.pdf> through edge-based algorithm <http://www.ngds-ku.org/pub/confabst0.htm#C42>. The procedure was repeated to link first phase to the second phase and so on. To construct a vertebral-level model, spinal column was visualized as a crystal structure <http://www.ngds-ku.org/Presentations/BZU1.pdf>. The center-of-mass of each vertebra was expressed in terms of positional coordinates in the body-coordinate system, just like *form factor* in crystallography. The surface structure of each vertebra could be studied using moiré fringe topography, which gives 3-D-coordinate information <http://www.ngds-ku.org/Presentations/Moire.pdf> and rasterstereography, which gives curvature information <http://www.ngds-ku.org/Presentations/Rasterstereography.pdf>. Combined with rotational (in terms of Euler angles) as well as inter-vertebral-spacing information, such an analysis may be considered equivalent to *structure factor*, determined by crystallographers. In order to study the vertebrae of spinal column of a live human being (without cutting open to reach vertebral surface to study using stereophotogrammetric techniques), *bone-surface-structure scanning* was introduced on May 30, 2013 <http://www.ngds-ku.org/Presentations/Backscatter.pdf> by the first author and elaborated in the Syed Firdous memorial lecture <http://www.ngds-ku.org/Presentations/Scan.pdf>, expanding upon the backscatter-body-scanning system, employed for passenger security screening at airports <http://www.ngds-ku.org/Presentations/Security.pdf>. Increasing the intensity of backscatter-X rays enough to reach to the bone surface and projecting these X rays through a moiré <http://www.ngds-ku.org/Papers/C68.pdf> or a dotted-raster grid <http://www.ngds-ku.org/Papers/J31.pdf>, vertebral surfaces could be studied by generating their height and curvature maps. In *grating-hologram-type-backscatter moiré* passive (post-recorded) correction (for misalignment or rotation) was possible. A projected-dotted-raster grid (X rays blocking raster patterns) generated 3-D-curvature map of the spinal column. Holographic images could be generated by recording and processing multiple images in the semi-circular angular range. As regards radiation damage, backscatter-X rays reflected off the bone surface and were not absorbed in its material, it was assumed that there would not be so much damage as from regular-X rays, provided eyes and gonads are, properly, shielded. Discretion is advised, while using these techniques in high-risk groups. Using techniques from physics (image projection/recording), computer science (edge-based and intensity-based algorithms) and information technology (image processing), stereophotogrammetric imaging of humans using moiré fringe topography, dotted-rasterstereography, raster stereophotogrammetry <http://www.ngds-ku.org/Presentations/Anthroimaging.pdf>, edge-based moiré and edge-based dotted-raster have the potential to open up new horizons in 3-D imaging of human-skin, -bone and -organ surfaces (sectional-semicircular views) as well as their infinitesimal movements.

**Keywords:** Human spinal column, static model, dynamic model, crystal-structure-based model, moiré fringe topography, dotted-rasterstereography, backscatter-X-ray scanning

**Conflict of Interest Statement:** No potential conflict of interest is identified for this work

**Web address of this document:** <http://www.ngds-ku.org/Presentations/Spinal.pdf>

**HTML version:** <http://www.ngds-ku.org/pub/confabstB.htm#C123>:

\*PhD (Mathematical Neuroscience); MSc, BSc (Honors), *both summa cum laude*, University of Karachi; MA, Johns Hopkins, Baltimore, MD, United States; MS, Indiana, Bloomington, IN, United States; Project Director, the NGDS Pilot Project; Director, SF Growth-and-Imaging Laboratory; Sessional Faculty, the Aga Khan University Medical College (1996-2006); Associated Professor in Orthopedic Surgery, Malmö General Hospital, Sweden (1988); Research Associate in Orthopedic Surgery, James Whitecomb Riley Hospital for Children, Indianapolis, IN, United States (1980); Summer Luncheon Seminar, 3-D Optical Imaging (for detection of scoliosis), Harvard-MIT Center for Biomedical Engineering, Massachusetts Institute of Technology, Cambridge, MA, United States and Member, Subject Committee for Physical Education, Health and Sport Sciences, National Testing Service, Islamabad, Pakistan; *paper mail:* Professor and Chairman, Department of Mathematics, University of Karachi, Karachi 75270, Pakistan; *telephone:* +92 21 9926 1300-15 ext. 2293; *homepage:* <http://www.ngds-ku.org/kamal>; *project URL:* <http://ngds-ku.org>